

Alcohol Symptom Checklist			Other Drugs Symptom Checklist		
In the past three months, have you:			In the past three months, have you:		
1. Had times when you ended up drinking more, or for longer than you intended?	Y	N	1. Had times when you ended up using drugs more, or for longer than you intended?	Y	N
2. More than once, wanted to cut down or stop drinking, or tried to, but couldn't?	Y	N	2. More than once, wanted to cut down or stop using drugs, or tried to, but couldn't?	Y	N
3. Spent a lot of time drinking, being sick after drinking, or getting over the after-effects?	Y	N	3. Spent a lot of time using drugs, being sick after use, or getting over the after-effects?	Y	N
4. Experienced craving — a strong need, or urge, to drink?	Y	N	4. Experienced craving — a strong need, or urge, to use drugs?	Y	N
5. Found that drinking — or being sick from drinking — often interfered with taking care of your home or family, caused job troubles or school problems?	Y	N	5. Found that using drugs — or being sick from using drugs — often interfered with taking care of your home or family, caused job troubles or school problems?	Y	N
6. Continued to drink even though it was causing trouble with your family or friends?	Y	N	6. Continued to use drugs even though it was causing trouble with your family or friends?	Y	N
7. Given up or cut back on activities that were important or interesting to you, or gave you pleasure, in order to drink?	Y	N	7. Given up or cut back on activities that were important or interesting to you, or gave you pleasure, in order to use drugs?	Y	N
8. More than once, gotten into situations while or after drinking that increased your chances of getting hurt (such as driving, swimming, using machinery, walking in a dangerous area or having unsafe sex)?	Y	N	8. More than once, gotten into situations while or after using drugs that increased your chances of getting hurt (such as driving, swimming, using machinery, walking in a dangerous area or having unsafe sex)?	Y	N
9. Continued to drink even though it was making you feel depressed or anxious or adding to another health problem, or after having had a memory blackout?	Y	N	9. Continued to use drugs even though it was making you feel depressed or anxious or adding to another health problem, or after having had a memory blackout?	Y	N
10. Had to drink much more than you once did to get the effect you want, or found that your usual number of drinks had much less effect than before?	Y	N	10. Had to use drugs much more than you once did to get the effect you want, or found that your usual number of drinks had much less effect than before?	Y	N
11. Found that when the effects of alcohol were wearing off, you had withdrawal symptoms, such as trouble sleeping, shakiness, irritability, anxiety, depression, restlessness, nausea or sweating, or sensed things that were not there?	Y	N	11. Found that when the effects of drugs were wearing off, you had withdrawal symptoms, such as trouble sleeping, shakiness, irritability, anxiety, depression, restlessness, nausea or sweating, or sensed things that were not there?	Y	N
<b>TOTAL:</b>			<b>TOTAL:</b>		

### Interpreting Symptom Checklist Results

2-3 symptoms indicate mild alcohol and/or other drug use disorder.

4-5 symptoms indicate moderate alcohol and/or other drug use disorder.

6+ symptoms indicate severe alcohol and/or other drug use disorder.